




## *Herpestes sanguineus*, Common Slender Mongoose

Assessment by: Do Linh San, E. & Maddock, A.H.



*View on [www.iucnredlist.org](http://www.iucnredlist.org)*

**Citation:** Do Linh San, E. & Maddock, A.H. 2016. *Herpestes sanguineus*. The IUCN Red List of Threatened Species 2016: e.T41606A45206143. <http://dx.doi.org/10.2305/IUCN.UK.2016-1.RLTS.T41606A45206143.en>

**Copyright:** © 2016 International Union for Conservation of Nature and Natural Resources

*Reproduction of this publication for educational or other non-commercial purposes is authorized without prior written permission from the copyright holder provided the source is fully acknowledged.*

*Reproduction of this publication for resale, reposting or other commercial purposes is prohibited without prior written permission from the copyright holder. For further details see [Terms of Use](#).*

*The IUCN Red List of Threatened Species™ is produced and managed by the [IUCN Global Species Programme](#), the [IUCN Species Survival Commission](#) (SSC) and [The IUCN Red List Partnership](#). The IUCN Red List Partners are: [BirdLife International](#); [Botanic Gardens Conservation International](#); [Conservation International](#); [Microsoft](#); [NatureServe](#); [Royal Botanic Gardens, Kew](#); [Sapienza University of Rome](#); [Texas A&M University](#); [Wildscreen](#); and [Zoological Society of London](#).*

*If you see any errors or have any questions or suggestions on what is shown in this document, please provide us with [feedback](#) so that we can correct or extend the information provided.*

## Taxonomy

Kingdom	Phylum	Class	Order	Family
Animalia	Chordata	Mammalia	Carnivora	Herpestidae

**Taxon Name:** *Herpestes sanguineus* (Rüppell, 1835)

**Synonym(s):**

- *Galerella sanguinea* (Rüppell, 1836)

**Common Name(s):**

- English: Common Slender Mongoose, Slender Mongoose
- French: Mangouste rouge

**Taxonomic Notes:**

In accordance with Hoffman and Taylor (2013) this assessment does not include *Herpestes ochraceus*, which is treated separately following many sources, notably Taylor (2013). Sometimes included in the genus *Galerella* (e.g., Wozencraft 1993, 2005; Veron *et al.* in press).

## Assessment Information

**Red List Category & Criteria:** Least Concern [ver 3.1](#)

**Year Published:** 2016

**Date Assessed:** February 28, 2015

**Justification:**

Listed as Least Concern as the species is common and widespread in a variety of habitats, there are no major threats, and it is present in several protected areas across the range.

**Previously Published Red List Assessments**

2008 – Least Concern (LC) – <http://dx.doi.org/10.2305/IUCN.UK.2008.RLTS.T41606A10505689.en>

1996 – Lower Risk/least concern (LR/lc)

## Geographic Range

**Range Description:**

One of the most widely distributed African mongooses, ranging from Senegal in the west to the Red Sea coast in Sudan in the east and south to the Northern Cape in South Africa (Hoffmann and Taylor 2013). Bahaa-el-din *et al.* (2013) recently obtained the first records of Common Slender Mongoose in Gabon, >350 km outside its previous known range. Past records of this species on the Cape Verde archipelago are in error (Masseti 2010, Hazevoet and Masseti 2011) and are in fact based on confusion with occurrence on the mainland Cape Verde itself. Stuart (1981) mentions a museum record of this species from Mountain Zebra N.P., but this specimen is not mentioned in the studies of Watson and Dippenaar (1987) and Watson (1990), and their most southerly distribution limit is probably the far eastern part of

the Eastern Cape in South Africa (Hoffmann and Taylor 2013). Also occurs on Zanzibar (Stuart and Stuart 1998, Goldman and Winther-Hansen 2003). Ranges from sea level to 2,700 m a.s.l. in the Ethiopian Highlands (Yalden *et al.* 1996).

### **Country Occurrence:**

**Native:** Angola (Angola); Benin; Botswana; Burkina Faso; Cameroon; Central African Republic; Chad; Congo; Congo, The Democratic Republic of the; Côte d'Ivoire; Djibouti; Equatorial Guinea; Eritrea; Ethiopia; Gabon; Gambia; Ghana; Guinea; Guinea-Bissau; Kenya; Liberia; Malawi; Mauritania; Mozambique; Namibia; Niger; Nigeria; Rwanda; Senegal; Sierra Leone; Somalia; South Africa; Sudan; Swaziland; Tanzania, United Republic of; Togo; Uganda; Zambia; Zimbabwe

# Distribution Map

*Herpestes sanguineus*




## Range

Extant (resident)

## Compiled by:

IUCN (International Union for Conservation of Nature)

NE DD LC > NT VU EN CR EW EX  
LEAST CONCERN



## **Population**

Common Slender Mongooses are among the most common mongooses in Africa. In the Serengeti N. P. (Tanzania), population densities between 1975 and 1990 ranged from 3–6 individuals/km<sup>2</sup> (Waser *et al.* 1995).

**Current Population Trend:** Stable

## **Habitat and Ecology (see Appendix for additional information)**

Present in a wide variety of habitats, but absent from true deserts and in sub-desertic parts of the Sahara such as Air, Niger. They occur on forest fringes, and may penetrate into forests along roads and are sometimes found around villages (Hoffmann and Taylor 2013). Slender Mongooses are generalist carnivores, their diet primarily comprising small vertebrates and invertebrates (Maddock 1988, Hoffmann and Taylor 2013).

**Systems:** Terrestrial

## **Use and Trade**

They have been recorded in bushmeat markets (e.g., Colyn *et al.* 2004) and Cunningham and Zondi (1991) listed this species among those used in traditional medicine in KwaZulu-Natal, South Africa.

## **Threats (see Appendix for additional information)**

There are no major threats to the species.

## **Conservation Actions (see Appendix for additional information)**

Present in numerous protected areas across their range.

## **Credits**

**Assessor(s):** Do Linh San, E. & Maddock, A.H.

**Reviewer(s):** Duckworth, J.W. & Hoffmann, M.

**Contributor(s):** Hoffmann, M. & Pacifici, M.

## Bibliography

- Bahaa-el-din, L., Henschel, P., Aba'a, R., Abernethy, K., Bohm, T., Bout, N., Coad, L., Head, J., Inoue, E., Lahm, S., Lee, M. E., Maisels, F., Rabanal, L., Starkey, M., Taylor, G., Vanthomme, A., Nakashima, Y. and Hunter, L. 2013. Notes on the distribution and status of small carnivores in Gabon. *Small Carnivore Conservation* 48: 19-29.
- Colyn, M., Dufour, S., Condé, P.C. and Van Rompaey, H. 2004. The importance of small carnivores in forest bushmeat hunting in the Classified Forest of Diecké, Guinea. *Small Carnivore Conservation* 31: 15-18.
- Cunningham, A.B. and Zondi, A.S. 1991. Use of animal parts for the commercial trade in traditional medicines. Institute of Natural Resources, University of Natal, Pietermaritzburg, South Africa.
- Goldman, H.V. and Winther-Hansen, J. 2003. First photographs of the Zanzibar servaline Genet *Genetta servalina archeri* and other endemic subspecies on the island of Unguja, Tanzania. *Small Carnivore Conservation* 29: 1-4.
- Hazevoet, C.J. and Masseti, M. 2011. On the history of the Green Monkey *Chlorocebus sabaeus* (L., 1766) in the Cape Verde Islands, with notes on other introduced mammals. *Zoologia Caboverdiana* 2: 12-24.
- Hoffmann, M. and Taylor, M.E. 2013. *Herpestes sanguineus* Slender Mongoose. In: J. Kingdon and M. Hoffmann (eds), *The Mammals of Africa. V. Carnivores, Pangolins, Equids and Rhinoceroses*, pp. 314-319. Bloomsbury, London, UK.
- IUCN. 2016. The IUCN Red List of Threatened Species. Version 2016-1. Available at: [www.iucnredlist.org](http://www.iucnredlist.org). (Accessed: 30 June 2016).
- Maddock, A.H. 1988. Resource partitioning in a viverrid assemblage. Ph.D. Thesis, University of Natal.
- Masseti, M. 2010. Mammals of the Macaronesian islands (the Azores, Madeira, the Canary and Cape Verde islands): redefinition of the ecological equilibrium. *Mammalia* 74: 3-34.
- Stuart, C. and Stuart, T. 1998. A note on the herpestids and viverrids of south-eastern Unguja (Zanzibar) Island. *Small Carnivore Conservation* 18: 16-17.
- Stuart, C.T. 1981. Notes on the mammalian carnivores of the Cape Province, South Africa. *Bontebok* 1: 1-58.
- Taylor, M.E. 2013. *Herpestes ochraceus* Somali Slender Mongoose. In: J. Kingdon and M. Hoffmann (eds), *The Mammals of Africa. V. Carnivores, Pangolins, Equids and Rhinoceroses*, pp. 310-311. Bloomsbury, London, UK.
- Veron, G., Patou, M.-L. and Jennings, A.P. In press. Systematics and evolution of the mongooses (Herpestidae, Carnivora). In: E. Do Linh San, J.J. Sato, J.L. Belant and M.J. Somers (eds), *Small Carnivores: Evolution, Ecology, Behaviour and Conservation*. Wiley-Blackwell, Oxford, UK.
- Waser, P.M., Elliott, L.F., Creel, N.M. and Creel, S.R. 1995. Habitat variation and mongoose demography. In: A.R.E. Sinclair and P. Arcese (eds), *Serengeti II: dynamics, management, and conservation of an ecosystem*, pp. 421-447. University of Chicago Press, Chicago, IL, USA.
- Watson, J.P. 1990. The taxonomic status of the slender mongoose, *Galerella sanguinea* (Ruppell, 1836) in southern Africa. *Navorsinge van die Nasionale Museum, Bloemfontein* 6: 351-492.
- Watson, J.P. and Dippenaar, N.J. 1987. The species limits of *Galerella sanguinea* (Ruppell, 1836), G.

*pulverulenta* (Wagner, 1839) and *G. nigrata* (Thomas, 1928) in southern Africa (Carnivora: Viverridae). *Navorsinge van die Nasionale Museum, Bloemfontein* 5: 351-492.

Wozencraft, W.C. 1993. Order Carnivora. In: D.E. Wilson and D.M. Reeder (eds), *Mammal Species of the World: A Taxonomic and Geographic Reference. Second Edition*, pp. 279-344. Smithsonian Institution Press, Washington, DC, USA.

Wozencraft, W.C. 2005. Order Carnivora. In: D.E. Wilson and D.M. Reeder (eds), *Mammal Species of the World: A Taxonomic and Geographic Reference. Third Edition*, pp. 532-628. Johns Hopkins University Press, Baltimore.

Yalden, D.W., Largen, M.J., Kock, D. and Hillman, J.C. 1996. Catalogue of the Mammals of Ethiopia and Eritrea. 7. Revised checklist, zoogeography and conservation. *Tropical Zoology* 9(1): 73-164.

## Citation

Do Linh San, E. & Maddock, A.H. 2016. *Herpestes sanguineus*. *The IUCN Red List of Threatened Species* 2016: e.T41606A45206143. <http://dx.doi.org/10.2305/IUCN.UK.2016-1.RLTS.T41606A45206143.en>

## Disclaimer

To make use of this information, please check the [Terms of Use](#).

## External Resources

For [Images and External Links to Additional Information, please see the Red List website](#).

## Appendix

### Habitats

(<http://www.iucnredlist.org/technical-documents/classification-schemes>)

Habitat	Season	Suitability	Major Importance?
1. Forest -> 1.5. Forest - Subtropical/Tropical Dry	-	Suitable	-
1. Forest -> 1.6. Forest - Subtropical/Tropical Moist Lowland	-	Suitable	-
2. Savanna -> 2.1. Savanna - Dry	-	Suitable	-
3. Shrubland -> 3.5. Shrubland - Subtropical/Tropical Dry	-	Suitable	-
3. Shrubland -> 3.7. Shrubland - Subtropical/Tropical High Altitude	-	Suitable	-
4. Grassland -> 4.5. Grassland - Subtropical/Tropical Dry	-	Suitable	-
14. Artificial/Terrestrial -> 14.4. Artificial/Terrestrial - Rural Gardens	-	Marginal	-

### Threats

(<http://www.iucnredlist.org/technical-documents/classification-schemes>)

Threat	Timing	Scope	Severity	Impact Score
5. Biological resource use -> 5.1. Hunting & trapping terrestrial animals -> 5.1.1. Intentional use (species is the target)	Ongoing	-	No decline	-
Stresses: 2. Species Stresses -> 2.1. Species mortality				

### Conservation Actions in Place

(<http://www.iucnredlist.org/technical-documents/classification-schemes>)

Conservation Actions in Place
In-Place Land/Water Protection and Management
Occur in at least one PA: Yes

### Additional Data Fields

Distribution
Lower elevation limit (m): 0
Upper elevation limit (m): 2700
Population
Population severely fragmented: No

**Habitats and Ecology**

Generation Length (years): 3

## The IUCN Red List Partnership



The IUCN Red List of Threatened Species™ is produced and managed by the [IUCN Global Species Programme](#), the [IUCN Species Survival Commission \(SSC\)](#) and [The IUCN Red List Partnership](#).

The IUCN Red List Partners are: [BirdLife International](#); [Botanic Gardens Conservation International](#); [Conservation International](#); [Microsoft](#); [NatureServe](#); [Royal Botanic Gardens, Kew](#); [Sapienza University of Rome](#); [Texas A&M University](#); [Wildscreen](#); and [Zoological Society of London](#).