


## *Atilax paludinosus*, Marsh Mongoose

Assessment by: Do Linh San, E., Angelici, F.M., Maddock, A.H., Baker, C.M. & Ray, J.



View on [www.iucnredlist.org](http://www.iucnredlist.org)

**Citation:** Do Linh San, E., Angelici, F.M., Maddock, A.H., Baker, C.M. & Ray, J. 2015. *Atilax paludinosus*. The IUCN Red List of Threatened Species 2015: e.T41590A45204865.  
<http://dx.doi.org/10.2305/IUCN.UK.2015-4.RLTS.T41590A45204865.en>

**Copyright:** © 2015 International Union for Conservation of Nature and Natural Resources

Reproduction of this publication for educational or other non-commercial purposes is authorized without prior written permission from the copyright holder provided the source is fully acknowledged.

Reproduction of this publication for resale, reposting or other commercial purposes is prohibited without prior written permission from the copyright holder. For further details see [Terms of Use](#).

The IUCN Red List of Threatened Species™ is produced and managed by the [IUCN Global Species Programme](#), the [IUCN Species Survival Commission \(SSC\)](#) and [The IUCN Red List Partnership](#). The IUCN Red List Partners are: [BirdLife International](#); [Botanic Gardens Conservation International](#); [Conservation International](#); [Microsoft](#); [NatureServe](#); [Royal Botanic Gardens, Kew](#); [Sapienza University of Rome](#); [Texas A&M University](#); [Wildscreen](#); and [Zoological Society of London](#).

If you see any errors or have any questions or suggestions on what is shown in this document, please provide us with [feedback](#) so that we can correct or extend the information provided.

## Taxonomy

Kingdom	Phylum	Class	Order	Family
Animalia	Chordata	Mammalia	Carnivora	Herpestidae

**Taxon Name:** *Atilax paludinosus* (G.[Baron] Cuvier, 1829)

### Common Name(s):

- English: Marsh Mongoose, Water Mongoose
- French: Mangouste des marais

## Assessment Information

**Red List Category & Criteria:** Least Concern [ver 3.1](#)

**Year Published:** 2015

**Date Assessed:** February 28, 2015

### Justification:

This species is listed as Least Concern because it has a wide distribution range, is generally common where there is suitable habitat, and occurs in several protected areas. There is no reason to believe that the species is declining at a rate fast enough to warrant listing in a threat category, and it does not meet any of the other IUCN Red List Criteria for even Near Threatened.

### Previously Published Red List Assessments

2008 – Least Concern (LC) – <http://dx.doi.org/10.2305/IUCN.UK.2008.RLTS.T41590A10487772.en>

1996 – Lower Risk/least concern (LR/lc)

## Geographic Range

### Range Description:

This species is widely distributed from Senegal, Guinea-Bissau and Sierra Leone eastward to southern Sudan and Ethiopia and south to southern Africa, where it is absent from most of Namibia, Botswana and large parts of central South Africa, wherever adequate water and cover are unavailable (Baker and Ray 2013). It is present on Pemba Island, but absent from Zanzibar (Pakenham 1984). It has been recorded from sea level to altitudes of 3,950 m asl in Bale Mountains N.P., Ethiopia (Yalden *et al.* 1996). Wozencraft (2005) lists this species as occurring in Algeria, presumably an error as there are no confirmed records from this country (M. Ahmim pers. comm. 2013).

### Country Occurrence:

**Native:** Angola (Angola); Benin; Botswana; Burkina Faso; Burundi; Cameroon; Central African Republic; Chad; Congo; Congo, The Democratic Republic of the; Côte d'Ivoire; Equatorial Guinea; Ethiopia; Gabon; Gambia; Ghana; Guinea; Guinea-Bissau; Kenya; Lesotho; Liberia; Malawi; Mozambique; Namibia; Niger; Nigeria; Rwanda; Senegal; Sierra Leone; Somalia; South Africa; Sudan; Swaziland; Tanzania, United Republic of; Togo; Uganda; Zambia; Zimbabwe

## Distribution Map




### *Atilax paludinosus*

#### Range

Extant (resident)

Compiled by:  
IUCN (International Union for  
Conservation of Nature)



The boundaries and names shown and the designations used on this map do not imply any official endorsement, acceptance or opinion by IUCN.



## Population

It is generally common in suitable habitat. It was the second most photographed species in a camera-trapping study in the Udzungwa Mountains, Tanzania (De Luca and Mpunga 2005). In KwaZulu-Natal, South Africa, the density was recorded at 1.8 individuals/km<sup>2</sup> (Maddock 1988).

**Current Population Trend:** Decreasing

## Habitat and Ecology (see Appendix for additional information)

It is mainly restricted to riparian habitats (rivers, streams, swamps, marshes and dams), wherever there is suitable vegetation cover and water in close proximity. It may also be found along estuaries and in coastal areas. Sometimes it is found away from watercourses, though only for limited periods (Baker and Ray 2013). Diet comprises mostly aquatic prey with crustaceans often dominating, which is unusual among mongooses (Baker and Ray 2013, Do Linh San *et al.* unpublished data). However, in some areas rodents constitute the main prey in terms of biomass (Maddock 1988).

**Systems:** Terrestrial, Freshwater

## Use and Trade

The Marsh Mongoose is commonly found in bushmeat markets. It was the most common carnivore appearing in bushmeat markets in southeast Nigeria (Angelici *et al.* 1999) and was also commonly recorded in bushmeat surveys in the Classified Forest of Diecke, Guinea (Colyn *et al.* 2004).

## Threats (see Appendix for additional information)

There are currently no major threats known to the species although hunting, habitat loss and encroachment by humans into its habitat are likely reasons for its declining population. Since it is dependent on riverine vegetation for shelter, the loss of this habitat may result in some localised declines where habitat loss is taking place (Baker and Ray 2013). The effects of water pollution on local availability of food resources and therefore Marsh Mongoose populations is unknown and would warrant investigation. The drainage of swamplands for conversion to arable land has been identified as a threat to this species in eastern Africa (Andama 2000).

## Conservation Actions (see Appendix for additional information)

This species is present in several protected areas across its range.

## Credits

**Assessor(s):** Do Linh San, E., Angelici, F.M., Maddock, A.H., Baker, C.M. & Ray, J.

**Reviewer(s):** Duckworth, J.W. & Hoffmann, M.

**Contributor(s):** Hoffmann, M.

## Bibliography

Andama, E. 2000. Status and Distribution of Carnivores in Bwindi Impenetrable National Park Southwestern Uganda. Makerere University, Uganda.

Angelici, F.M., Luiselli, L., Politano, E. and Akani, G.C. 1999. Bushmen and mammal fauna: A survey of the mammals traded in bush-meat markets of local people in the rainforests of southeastern Nigeria. *Anthropozoologica* 30: 51-58.

Baker, C.M. and Ray, J.C. 2013. *Atilax paludinosus* Marsh Mongoose. In: J. Kingdon and M. Hoffmann (eds), *The Mammals of Africa. V. Carnivores, Pangolins, Equids and Rhinoceroses*, pp. 298-302. Bloomsbury, London, UK.

Colyn, M., Dufour, S., Condé, P.C. and Van Rompaey, H. 2004. The importance of small carnivores in forest bushmeat hunting in the Classified Forest of Diecké, Guinea. *Small Carnivore Conservation* 31: 15-18.

De Luca, D.W. and Mpunga, N.E. 2005. Carnivores of the Udzungwa Mountains: Presence, distributions and threats. Wildlife Conservation Society, Mbeya, Tanzania.

IUCN. 2015. The IUCN Red List of Threatened Species. Version 2015-4. Available at: [www.iucnredlist.org](http://www.iucnredlist.org). (Accessed: 19 November 2015).

Maddock, A.H. 1988. Resource partitioning in a viverrid assemblage. Ph.D. Thesis, University of Natal.

Pakenham, R.H.W. 1984. *The Mammals of Zanzibar and Pemba islands*. Printed Privately, Harpenden.

Wozencraft, W.C. 2005. Order Carnivora. In: D.E. Wilson and D.M. Reeder (eds), *Mammal Species of the World: A Taxonomic and Geographic Reference. Third Edition*, pp. 532-628. Smithsonian Institution Press, Washington, DC, USA.

Yalden, D.W., Largen, M.J., Kock, D. and Hillman, J.C. 1996. Catalogue of the Mammals of Ethiopia and Eritrea. 7. Revised checklist, zoogeography and conservation. *Tropical Zoology* 9(1): 73-164.

## Citation

Do Linh San, E., Angelici, F.M., Maddock, A.H., Baker, C.M. & Ray, J. 2015. *Atilax paludinosus*. *The IUCN Red List of Threatened Species 2015*: e.T41590A45204865. <http://dx.doi.org/10.2305/IUCN.UK.2015-4.RLTS.T41590A45204865.en>

## Disclaimer

To make use of this information, please check the [Terms of Use](#).

## External Resources

For [Images and External Links to Additional Information](#), please see the [Red List website](#).

# Appendix

## Habitats

(<http://www.iucnredlist.org/technical-documents/classification-schemes>)

Habitat	Season	Suitability	Major Importance?
1. Forest -> 1.8. Forest - Subtropical/Tropical Swamp	-	Suitable	Yes
4. Grassland -> 4.6. Grassland - Subtropical/Tropical Seasonally Wet/Flooded	-	Marginal	-
5. Wetlands (inland) -> 5.1. Wetlands (inland) - Permanent Rivers/Streams/Creeks (includes waterfalls)	-	Suitable	Yes
5. Wetlands (inland) -> 5.2. Wetlands (inland) - Seasonal/Intermittent/Irregular Rivers/Streams/Creeks	-	Suitable	Yes
5. Wetlands (inland) -> 5.3. Wetlands (inland) - Shrub Dominated Wetlands	-	Suitable	Yes
5. Wetlands (inland) -> 5.4. Wetlands (inland) - Bogs, Marshes, Swamps, Fens, Peatlands	-	Suitable	Yes
5. Wetlands (inland) -> 5.7. Wetlands (inland) - Permanent Freshwater Marshes/Pools (under 8ha)	-	Suitable	Yes
9. Marine Neritic -> 9.10. Marine Neritic - Estuaries	-	Suitable	Yes
13. Marine Coastal/Supratidal -> 13.5. Marine Coastal/Supratidal - Coastal Freshwater Lakes	-	Suitable	Yes
15. Artificial/Aquatic & Marine -> 15.1. Artificial/Aquatic - Water Storage Areas (over 8ha)	-	Suitable	Yes

## Threats

(<http://www.iucnredlist.org/technical-documents/classification-schemes>)

Threat	Timing	Scope	Severity	Impact Score
1. Residential & commercial development -> 1.1. Housing & urban areas	Ongoing	Minority (50%)	Negligible declines	Low impact: 4
	Stresses:	1. Ecosystem stresses -> 1.1. Ecosystem conversion 1. Ecosystem stresses -> 1.2. Ecosystem degradation 2. Species Stresses -> 2.2. Species disturbance		
1. Residential & commercial development -> 1.2. Commercial & industrial areas	Ongoing	Minority (50%)	Negligible declines	Low impact: 4
	Stresses:	1. Ecosystem stresses -> 1.1. Ecosystem conversion 1. Ecosystem stresses -> 1.2. Ecosystem degradation 2. Species Stresses -> 2.2. Species disturbance		
5. Biological resource use -> 5.1. Hunting & trapping terrestrial animals -> 5.1.1. Intentional use (species is the target)	Ongoing	Minority (50%)	Slow, significant declines	Low impact: 5
	Stresses:	2. Species Stresses -> 2.1. Species mortality		

7. Natural system modifications -> 7.2. Dams & water management/use -> 7.2.7. Abstraction of ground water (agricultural use)	Ongoing	Minority (50%)	Slow, significant declines	Low impact: 5
Stresses:		1. Ecosystem stresses -> 1.1. Ecosystem conversion 1. Ecosystem stresses -> 1.2. Ecosystem degradation		

## Conservation Actions in Place

(<http://www.iucnredlist.org/technical-documents/classification-schemes>)

<b>Conservation Actions in Place</b>
In-Place Land/Water Protection and Management
Occur in at least one PA: Yes

## Research Needed

(<http://www.iucnredlist.org/technical-documents/classification-schemes>)

<b>Research Needed</b>
1. Research -> 1.2. Population size, distribution & trends
1. Research -> 1.5. Threats

## Additional Data Fields

<b>Distribution</b>
Lower elevation limit (m): 0
Upper elevation limit (m): 3950
<b>Population</b>
Population severely fragmented: No
<b>Habitats and Ecology</b>
Generation Length (years): 4

## The IUCN Red List Partnership



The IUCN Red List of Threatened Species™ is produced and managed by the [IUCN Global Species Programme](#), the [IUCN Species Survival Commission \(SSC\)](#) and [The IUCN Red List Partnership](#).

The IUCN Red List Partners are: [BirdLife International](#); [Botanic Gardens Conservation International](#); [Conservation International](#); [Microsoft](#); [NatureServe](#); [Royal Botanic Gardens, Kew](#); [Sapienza University of Rome](#); [Texas A&M University](#); [Wildscreen](#); and [Zoological Society of London](#).